

THE RECOMMENDED VACCINE SCHEDULE FROM BIRTH TO RETIREMENT



- Respiratory syncytial virus (RSV-mAb [Nirsevimab])
- Hepatitis B (HepB)
- Rotavirus (RV) (RV1 (2-dose series); RV5 (3-dose series))
- Diphtheria, tetanus, & acellular pertussis (DTaP: <7 yrs)
- Haemophilus influenzae type b (Hib)
- Pneumococcal conjugate (PCV15, PCV20)
- Inactivated poliovirus (IPV: <18 yrs)
- COVID-19 (1vCOV-mRNA, 1vCOV-aPS) *
- Influenza (IIV4)
- Meningococcal (MenACWY-TT)
- Measles, mumps, rubella (MMR)
- Varicella (VAR)
- Hepatitis A (HepA)
- Tetanus, diphtheria, & acellular pertussis (Tdap) **
- Dengue (DEN4CYD: 9-16 yrs)
- Zoster recombinant (RZV)
- Human papillomavirus (HPV)
- Pneumococcal poly-saccharide vaccine (PPSV23)

Important Note: The vaccines are listed at the earliest ages recommended by the CDC, but the actual window may extend beyond that.

Annual vaccination



Footnotes:

- (a) Depending on maternal RSV vaccination status
- (b) If using a 3-dose series
- (c) Minimum 6 month interval
- (d) Begin age 9
- (e) Then Td or Tdap booster every 10 years
- (f) If born in 1957 or later

- (g) If born in 1980 or later
- (h) Depending on vaccine or condition
- (i) Depending on previous vaccination status
- (j) If no evidence of immunity
- (k) 75 and older

This is the vaccine schedule recommended by the CDC for everyone. There may be special circumstances or conditions that affect when a vaccine should be administered or who should receive it. Please refer to these complete vaccine schedules provided by the CDC to learn about these unique situations, and as always, consult with your doctor:

Child and Adolescent Immunization Schedule by Age:
<https://www.cdc.gov/vaccines/hcp/imz-schcdulcs/child-adolescscnt-agc.html>

Adult Immunization Schedule by Age:
<https://www.cdc.gov/vaccines/hcp/imz-schcdulcs/adult-agc.html>

* Recommended for all persons over 6 months. After the initial vaccination course is complete, it is recommended that those who are 75 years and older receive a booster every 6 months. For those who are immunocompromised, it is recommended that those who are 18-74 years old receive a booster every 12 months.
 ** 1 dose Tdap during each pregnancy and 1 dose Tdap for wound management (if vaccine history is unknown or previous vaccine was over 5 years ago).

SOURCES:

- www.cdc.gov/vaccines/schedules/index.html
- www.cdc.gov/vaccines/schedules/hcp/imz/child-adolescent.html
- www.cdc.gov/vaccines/schedules/hcp/imz/adult.html